

**Summary of Immunization Requirements for School, Daycare & Head Start**  
 (Refer to 902 KAR 2:060 effective July 1, 2011, for details on exceptions and exemptions)

If a child is this age	These immunizations are required for attendance
<3 months	None
≥3 - <5 months	1 DTaP or DTP; 1 IPV or OPV; 1 Hib; 1 Hep B; 1 PCV
≥5 - <7 months	2 DTaP or DTP, or combination of the two; 2 IPV or OPV, or combination of the two; 2 Hib; 2 Hep B; 2 PCV
≥7 - <12 months	3 DTaP or DTP, or combination of the two; 2 OPV or IPV, or combination of the two; 2 Hib <sup>1</sup> ; 2 Hep B; 3 PCV*
≥12 - <16 months	3 DTaP or DTP, or combination of the two; 2 OPV or IPV, or combination of the two; 3 Hib <sup>1,2</sup> ; 2 Hep B; 4 PCV*
≥16 - <19 months	4 DTaP or DTP, or a combination of the two; 2 OPV or IPV, or a combination of the two; 4 Hib <sup>1,2</sup> ; 2 Hep B; 4 PCV* <sup>3</sup> ; 1 MMR; 1 Varicella
≥19 - 48 months	4 DTaP or DTP, or a combination of the two; 3 OPV or IPV, or a combination of the two; 4 Hib <sup>1,2</sup> ; 3 Hep B; 4 PCV* <sup>8</sup> ; 1 MMR; 1 Varicella <sup>4</sup> ;
≥48 months - <5 years	4 DTaP or DTP, or a combination of the two; 3 OPV or IPV, or a combination of the two; 4 Hib <sup>1,2</sup> ; 3 Hep B; 4 PCV* <sup>3</sup> ; 2 MMR; 2 Varicella <sup>4</sup>
≥ 5 years - <7 years	5 DTaP or DTP <sup>5</sup> , or a combination of the two; 4 OPV or IPV <sup>6</sup> , or a combination of the two; 3 Hep B; 2 MMR; 2 Varicella <sup>4</sup>
≥7 years - <11 years	5 DTaP or DTP <sup>5</sup> , or combination of the two; OR 1 dose of Td that was preceded by (2) doses DTaP, DTP,DT Td or combination of the (4) vaccines; 4 OPV or IPV <sup>6</sup> , or combinations of the two; 3 HepB; 2 MMR; 2 Varicella
At 6 <sup>th</sup> grade entry	1 Tdap; 4 IPV or OPV or combinations of the 2 vaccines; 3 Hep B <sup>7</sup> ; 2 MMR; 2 Varicella; 1 MCV or MPSV.
≥10 years since last DTaP, DTP or Td	1 Tdap if no previous dose of Tdap has been administered. Otherwise, give one dose of Td.

**Notes:**

1. If first two doses of Hib vaccine were meningococcal protein conjugate, the third dose may be omitted and the child shall be considered as having received three (3) doses.
  2. If Hib vaccine has been administered on or after 15 months of age, the child is not required to have further doses.
  3. One dose of PCV vaccine must be administered on or after 12 months of age.
  4. Unless a physician states that the child has had chickenpox disease.
  5. If the 4<sup>th</sup> dose of DT, DTP or DTaP was administered on or after the fourth birthday, the fifth dose shall not be required.
  6. If the 3<sup>rd</sup> dose of IPV or OPV was administered on or after the fourth birthday, the fourth dose shall not be required if at least 6 months preceded the previous dose.
  7. Children 11-15 years of age may receive an alternate two-dose series consisting of two 10 mcg doses of Merck's RecombivaxHB separated by 4-6 months.
  8. If (1) dose of PCV has been administered on or after 24 months of age the child is not required to have further doses.
- \*A child who receives the first dose of PCV vaccine between 7 to 11 months of age may need fewer doses to complete the series. Refer to 902 KAR 2:060 and the ACIP Recommended Immunization Schedule for details.